



## Upper Endoscopy (EGD) Prep Guide

You have been scheduled for an Upper Endoscopy (EGD).

**Plan ahead to help reduce your stress.** Use these step-by-step instructions for a successful procedure so that your doctor can clearly view your stomach.

If you have any questions, please contact the Gastroenterology office at the Guthrie Clinic at (570) 887-2852.

Thank you for choosing Guthrie!

### Upper Endoscopy (EGD) Prep Guide

- **Health Considerations** page 2 **(review today)**
  - Medications and diagnoses
  - Patients on blood thinners
  - Diabetic patients
- **Get prepared** page 3 **(review 1-2 weeks before)**
  - Sedation & transportation requirements
- **Get started** page 4 **(review 5 days before the procedure)**
- **Procedure day reminders** page 4 **(review 5 days before & day of)**
- **Prep Instructions\*** **(follow the day before & day of)**
  - Morning procedure instructions page 5
  - Afternoon procedure instructions page 5

***\*Please note there are different instructions for scheduled appointments in the morning and scheduled appointments in the afternoon.***

## Health Considerations

(review today)

### Medications and diagnoses

Please call the Gastroenterology office at the Guthrie Clinic at (570) 887-2852, select option 1 **immediately** if **any** of the following apply to you:

- You use oxygen at home.
- You use CPAP or BIPAP or have been diagnosed with sleep apnea.
- You weigh over 300 lbs. or have a BMI (body mass index) over 45.
- If you are pregnant or think you may be pregnant.
- **If you take pain medications, anti-anxiety medications, psychiatric medications, use medical marijuana, or recreational drugs.**
- **If you need to cancel or reschedule your appointment.**

### Patients on blood thinners

If you use Aggrenox, Arixtra, Brilinta, Effient, Eliquis, Plavix, Pletal, Pradaxa, Ticlid, Xarelto, or any blood thinner (anticoagulant) or an anti-platelet drug, most patients need to stop taking these 5 days prior to procedure.

- Please speak with your physician who orders this medication before stopping.

If you use a blood thinner named Coumadin, Warfarin, or Jantoven you will need special instructions about stopping this drug before the procedure. ***The timing of this is very important!***

- If you are followed by the Coumadin Clinic, please contact the Sayre Coumadin Clinic at 570-887-2200 for instructions before stopping, to see if they prefer that you use an alternative anticoagulant.
- If you are NOT followed by the Coumadin Clinic, please contact your physician who orders this for instructions.

### Patient taking GLP-1 Receptor Agonists

- If you are taking medications such as Ozempic, Wegovy, Trulicity, Mounjaro, Saxenda, Rybelsus: Injectable form must be held for 1 week prior to your procedure and pill form must be held the day of the procedure.

## **Diabetic patients**

- If you take a pill to lower your sugar, do not take it on the day of your procedure.
- If you are taking regular Insulin® or NPH, do not take it on the day of your procedure.
- If you are taking any other insulin preparation such as Lantus, Humalog, or 70/30 insulin: It is recommended you take a ½ dose the evening prior to your procedure and no dose the day of your procedure.
- We recommend you have products with and without sugar on hand at home to keep your blood glucose in balance during the preparation period.

## **Get prepared**

(1-2 weeks before the procedure)

Clear Liquid Suggestions:

- Gatorade, Pedialyte, or Powerade
- Coffee or tea without creamer
- Carbonated and non-carbonated soft drinks
- Kool-Aid or other fruit-flavored drinks
- Apple juice, white cranberry juice, or white grape juice
- Jell-O, popsicles
- Chicken, beef, or vegetable broth

## **Sedation requires that patients make transportation arrangements**

Please be aware that you will be receiving sedation for your procedure. It will limit what you can do after your procedure until the following day.

### **ALL ENDOSCOPY PATIENTS MUST HAVE A DESIGNATED DRIVER**

**Due to sedation medication, it is unlawful to drive until the following day after a procedure and is considered driving under the influence of drugs (DUI), a punishable offense.**

**It is required that all endoscopy patients have a designated driver (18 years or older) available to drive the patient home after their procedure. You may not take public transportation unless accompanied by someone 18 years of age or older.**

## Get started

(5 days before the procedure)

### 5 days before your procedure, stop taking:

- **Iron/multivitamins:** Stop iron (such as ferrous sulfate) or vitamins containing iron.
- **Fiber supplements:** Stop fiber supplements such as Metamucil, Citrucel.
- **Blood Thinners:**
  - **If you use Aggrenox, Arixtra, Brilinta, Effient, Eliquis, Plavix, Pletal, Pradaxa, Ticlid, Xarelto, or any blood thinner (anticoagulant) or anti-platelet drug:** most patients need to stop taking these 5 days prior to procedure. **Please speak with your physician who orders this medication before stopping. See page 2.**
  - **If you use a blood thinner named Coumadin, Warfarin, or Jantoven you will need special instructions about stopping this drug before the procedure. See page 2.**

Please note that Aspirin or NSAID'S such as Advil, Ibuprofen, Aleve do NOT need to be stopped prior to your procedure.

## Procedure day reminders

(review 5 days prior, and day of)

### Medication on the day of your procedure

- **Do NOT take blood thinners, see page 2.**
- **Diabetics, see page 2.**
- **Do NOT take GLP-1 Receptor Agonists, see page 2.**
- You may take other medications (including pain medications, anti-anxiety medications, and blood pressure medications) with a **small sip of water**. Take inhaler as prescribed if needed in the morning.

### What to bring

- A list of your current medications, your insurance card, and **Photo ID**
- Hearing aids, inhaler, and/or ileostomy or colostomy supplies (as applicable)
- Wear comfortable, loose fitting clothing
- Please leave all valuables at home. Cell phones and tablets are not allowed after you enter the patient care area and must be turned off until after discharge. This is for the privacy of all our patients.

### Transportation & Discharge

- You must have a driver 18 years of age or older with a valid driver's license for this procedure.
- You may not take public transportation unless accompanied by someone 18 years of age or older.
- If someone is picking you up, we must be able to contact them by **phone before** your procedure.
- We will **not** continue with your procedure until we confirm with your driver by talking to them.
- You should plan on being at our facility for a minimum of **2-3 hours** for your appointment. Please plan accordingly, as **delays may occur**.

## **Upper Endoscopy (EGD) Prep Guide for Morning Procedures**

Follow these instructions if you are scheduled to arrive BEFORE Noon

### **The day before your procedure:**

- You are not allowed to eat or drink anything after midnight.

### **On the day of your procedure:**

- Review procedure day reminders, page 4.
- Do not take blood thinners the day of your procedure (see page 2).
- Do NOT take GLP-1 Receptor Agonists (see page 2).
- You may take other medications, with a small sip of water.
- You may brush your teeth.

## **Upper Endoscopy (EGD) Prep Guide for Afternoon Procedures**

Follow these instructions if you are scheduled to arrive AFTER Noon

### **The day before your procedure:**

- No solids foods after midnight.

### **On the day of your procedure:**

- Review procedure day reminders, page 4.
- You may have clear liquids only until 10:00 am.
- Do not take blood thinners the day of your procedure (see page 2).
- Do NOT take GLP-1 Receptor Agonists (see page 2).
- You may take other medications, with a small sip of water.
- You may brush your teeth.

